

Washington State
University

Oregon State
University

University of
Idaho



FERTILIZING WITH MANURE AND OTHER ORGANIC AMENDMENTS

PNW PEER
REVIEWED

PNW533

FERTILIZING WITH MANURE AND OTHER ORGANIC AMENDMENTS

Table of Contents

Quick Facts	3
Overview	4
Is Manure a Good Choice for Your Farm?	4
Local Sources of Manure	4
Quality of Manure Sources	4
Crop Rotations	4
Soil Fertility	4
Composted vs. Uncomposted Manure	4
How to Determine the Nutrient Content of Manure	5
Laboratory Analysis	5
Choosing a Laboratory	7
Sampling Manure	8
Published Values	8
Nitrogen Availability for Crop Growth	8
Calculating the Application Rate	10
Applying Manure	10
Manure Spreaders	10
Equipment	10
Calibrating Manure Applications	11
Timing Manure Applications	11
Using Manure Safely	11
Manure Storage	12
Using Soil Testing to Adjust Application Rates	12
Long-Term Effects of Manure Applications	13
Worksheet A	14
Spreader Calibration Using a Full Spreader Load and Scales	14
Worksheet B	15
Spreader Calibration Using Manure Bulk Density	15
Bulk Density Estimate	16
Worksheet C	17
Calculating Manure Application Rates	17
Additional Resources	18

By

Andy Bary, Senior Scientific Assistant, WSU Puyallup Research and Extension Center, **Craig Cogger**, Extension Soil Scientist Emeritus, WSU Puyallup Research and Extension Center, **Dan Sullivan**, Professor, Department of Crop and Soil Science, Oregon State University

Abstract

Are you thinking about using manure to fertilize your farm but want more information? Properly managed manure applications recycle nutrients to crops, improve soil quality, and protect water quality. From deciding whether manure is right for your farm to learning how to calibrate manure applications, this publication takes you through the process of fertilizing with manure and other organic amendments.

Fertilizing with Manure and Other Organic Amendments

Quick Facts

Manure nutrient management is a cyclic process that includes:

- determining crop nutrient need,
- choosing a nutrient source,
- selecting and calibrating a manure application rate to deliver a target nutrient rate,
- monitoring crop performance and soil test values, and
- adjusting future application rates based on crop and soil monitoring.

Manures are unbalanced in nutrient content. When applied to meet crop nitrogen need, they usually will supply more phosphorus and potassium than is necessary for crops. Monitoring phosphorus and potassium via soil testing every three to five years is recommended to avoid excessive accumulation.

Unlike fertilizer, manure has variable nutrient content, depending upon site-specific factors (e.g., bedding, moisture, and storage or composting). See Table 2 for typical values.

For the current growing season, apply manure based on its ability to supply plant-available nitrogen to meet, but not exceed crop needs. Keep in mind that over-application of some products can damage plants with too much soluble salt.

Composts are lower in plant-available nitrogen than are most raw manures. Composting raw manure kills weed seeds and pathogens, facilitates storage, and makes a more uniform material that is easier to apply. For typical compost nutrient analyses, see Table 3.

Byproducts from cities and food processing facilities have nutrient value and are sometimes available for use on farms. See Table 4 for examples.

Manures and specialty products that contain more than 4% total nitrogen should be applied at low rates based on plant-available nitrogen supplied. See Table 5.

Use published values for manure nutrient content (Tables 2–5) with caution. Laboratory analyses of manure can improve the accuracy of manure nutrient application rates. For best results, send samples to laboratories that have demonstrated proficiency in performing manure analyses. Under Additional Resource, see the Lists of Laboratories and Consultants section of this publication for more information. Other organic materials can be analyzed for nutrient content using the same laboratory methods used for manure.

Calibration worksheets for manure spreaders that provide several options for monitoring manure application rates are included in this publication. Keeping records of manure nutrient application rates on a field-by-field basis can assist in evaluating and modifying manure application programs.

This publication is written for both organic farmers and other farmers who use manure. For details on organic farming standards, refer to the *National Organic Standards* in Additional Resources.

Overview

Manure is a good source of plant nutrients and organic matter. Properly managed manure applications recycle nutrients to crops, improve soil quality, and protect water quality. Although the focus of this publication is on animal manures and manure composts, the same principles apply for other waste-derived organic amendments, such as peppermint hay (leaves and stems that have been heated to remove peppermint oil), fish waste, and composted yard debris and food waste.

Is Manure a Good Choice for Your Farm?

Consider the following questions before applying animal manure on your farm:

- Are there local sources of manure?
- What is the quality of local manure sources?
- Where does manure fit in your crop rotation?
- Will your crops respond to added nutrients?
- What are the environmental and food safety risks and how can you mitigate them?

Local Sources of Manure

If you raise livestock, or if you have neighbors who do, you will likely have plenty of manure available nearby. If you have to import manure from more distant sources, transportation can greatly increase costs. You will need to decide if it makes economic sense if you are considering importing manure. Other organic materials may also be locally available, such as peppermint hay in parts of the Willamette Valley and the Columbia Basin, or shrimp and crab waste if processing facilities are nearby.

Quality of Manure Sources

Manure quality varies considerably, as discussed later in this publication. You will need to compare the quality of available manure sources (nutrient supply, ease of handling, odor, and risk of weed seeds or pathogens) with your farming needs. See Table 1 for a comparison of composted and uncomposted manure sources.

Crop Rotations

Manure use is compatible with most crops and crop rotations. Remember that crop nutrient needs vary and manure applications should be consistent with the nutrient needs of your crops. Soil test results and Extension publications are good sources of information on crop nutrient needs.

Fresh manure is not a good choice for salad and root crops that are eaten raw because of food safety concerns. For more information, see the section titled Using Manure Safely.

Soil Fertility

Manure provides the most benefits on soils with deficient to adequate levels of nutrients. Soils with high to excessive levels of nutrients are not a good choice for manure use because the nutrients in manure are less likely to benefit crops and more likely to leach into groundwater or run off into surface water. If you have excessive levels of nutrients in your soil, limit applications of those nutrients to recommended starter levels and use an alternative source of organic matter input, such as growing cover crops.

Composted vs. Uncomposted Manure

Composting is more than just piling the material and letting it sit. Composting is the active management of manure and bedding to aid the decomposition of organic materials and destruction of pathogens by microorganisms under controlled conditions (Figure 1). For more information on composting, refer to the Additional Resources section.

What are the benefits and detriments of using uncomposted or composted manure? In some cases, they have comparable properties, and in other ways, they are quite different. Table 1 compares the two materials.



Figure 1. Composting manure on-farm using an aerated, static (untuned) pile. Photo by Andy Bary.

Table 1. Compost and manure comparison

Compost	Manure
Slow release of nutrients. Often best as a soil builder.	Usually higher nutrient content. Often used as a fertilizer.
Easier to spread.	Sometimes difficult to spread.
Lower potential to degrade water quality.	Higher potential to degrade water quality.
Less likely to contain weed seeds.	More likely to contain weed seeds.
Higher investment of time or money.	Lower investment of time or money.
Reduced pathogen levels (e.g., Salmonella and E. coli).	Potential for higher pathogen levels.
More expensive to purchase.	Less expensive to purchase.
Fewer odors (although poor composting conditions can create foul odors).	Odors sometimes.
Improves soil tilth.	Improves soil tilth.
More stabilized carbon; a greater fraction ends up as stable organic matter.	Less stabilized carbon; more is lost to microbial decomposition.

How to Determine the Nutrient Content of Manure

Not knowing the nutrient content of manure can lead to large errors in nutrient application rates. We strongly advise you test the manure you plan to use (Figure 2). If you buy manure from a commercial source, they should be able to provide you with nutrient test values; you would not need to do further testing. In the absence of test values, use the published values in Table 2 (manure) and Table 3 (manure composts) as a starting point. For other organic amendments, use the values in Table 4.



Figure 2. Nutrient concentrations and availability in manure varies widely depending on the type of livestock and how the manure is stored and handled. Photo by Andy Bary.

Organic Fertilizers vs. Soil Builders

Manure and other organic amendments that supply large amounts of nutrients are organic fertilizers. They are applied at rates estimated to meet crop nutrient needs. Some manures and amendments (including most composts) supply smaller amounts of nutrients and are used as soil builders. Farmers can apply these amendments at higher rates to build soil organic matter quickly without risk of over-applying nutrients.

Since manure application rates are usually based on nitrogen, much of the difference between organic fertilizers and soil builders depends on the concentration and availability of nitrogen. In general, manure that contains more than 2% nitrogen on a dry-weight basis is used as a fertilizer, while manure with less than 2% nitrogen has low enough nitrogen availability to be a soil builder. Most composts have low enough nitrogen availability to be used as soil builders. During composting, most of the easily decomposable or mineralizable forms of nitrogen are converted to more stable organic forms or lost as ammonia gas.

See the section on Nitrogen Availability for Crop Growth for background on release and availability of nitrogen from organic amendments. Tables 2–5 provide estimates of nitrogen concentration and availability from a variety of manures, composts, and other amendments.

Laboratory Analysis

Laboratory analysis measures the nutrients in manure instead of just estimating them based on table values. Using testing laboratories requires proper sampling techniques and timely delivery of the sample to the laboratory.

The main components you should have measured are total nitrogen, ammonium nitrogen, total phosphorus, total potassium, electrical conductivity, and dry matter. If the manure is old or has been composted, you may also want to test for nitrate-N. Electrical conductivity is a measure of soluble salts. It is an important factor in arid regions, where salts can accumulate in the soil and harm crop growth. Dry matter is used to convert lab results from a dry-weight to an as-is basis.

Table 2. Uncomposted animal manures: Typical nutrient content, dry matter, bulk density, and nitrogen availability^a

Type	N	P ^b	K ^b	Dry Matter	Density	Total N	C/N ratio	Available N ^c		
				lb/ton as-is ^d	%			lb/cubic yd ^e	% dry weight	% of total N
Broiler with litter	56	27	39	68	850	4.1	11	40 to 60	22–34	10–14
Laying hen	39	28	29	39	1400	5.1	8	40 to 60	16–24	11–16
Turkey	51	23	27	55	1000	4.7	9	40 to 60	20–30	10–15
Rabbit	20	11	12	35	1400	2.8	12	20 to 40	4–8	3–6
Sheep	17	7	15	32	1400	2.9	12	20 to 40	3–7	2–5
Goat	17	4	15	40	1400	2.2	14	15 to 30	2–5	2–4
Beef	14	4	12	28	1400	2.4	15	15 to 30	2–4	1.5–3
Llama	15	7	5	35	1400	2.1	15	15 to 30	2–4	1.5–3
Alpaca	13	8	5	27	1400	2.4	15	15 to 30	2–4	1.5–3
Stockpiled dairy manure ^f	18	6	35	50	1000	1.9	15	10 to 20	2–4	1–2
Horse no bedding	8	3	8	25	1400	1.6	20	0 to 15	0–1	0–0.7
Horse with bedding	10	3	10	35	1400	1.4	30	-5 ^g to 10	< 1	< 1
Dairy cow separated solids	6	1	3	21	1100	1.4	32	-5 to 10	< 1	< 1

^aData sources include Gale et al. 2006; Brown 2013; Bary 2010; and unpublished Pacific Northwest data from Bary, Cogger, and Sullivan, and A. Moore. See Additional Resources for citations.

^bManure analyses are usually reported as P and K, while fertilizer labels use P₂O₅ and K₂O. To convert from P to P₂O₅, multiply by 2.3. To convert from K to K₂O, multiply by 1.2.

^cAvailable N is estimated N that becomes available to plants during the first growing season after application.

^d“As-is” is typical dry matter content for solid manure stored under cover. Dry matter, nutrients, and density can vary widely depending on how manure was handled.

^eDensities listed as 1400 lb/cubic yard are estimates. Most manures and composts with high moisture (low dry matter) have a density close to 1400 lb/cubic yard. Use the dropped bucket method described in Worksheet B to get a more accurate estimate of bulk density.

^fStockpiled dairy manure data from Moore et al. 2015.

^gNegative value indicates nitrogen immobilization (conversion from available to unavailable form).

Table 3. Manure composts: Typical nutrient content, dry matter, bulk density, and nitrogen availability^a

	N	P ^b	K ^b	Dry Matter	Density	Total N	C/N ratio	Available N ^c		
				lb/ton as-is ^d	%			lb/cubic yd	% dry weight	% of total N
Broiler litter “compost” ^e	44	26	33	57	900	3.8	10	30 to 40	12–18	6–9
Rabbit manure compost	15	8	22	43	1300	1.8	10	15 to 30	2–5	1.5–3
Beef manure compost	18	7	22	64	900	1.4	11	0 to 10	2–4	0.8–1.6
Separated dairy solids compost	10	1	3	25	1400	2.1	18	0 to 10	0–1	0–0.7

^aData sources include Gale et al. 2006; Larney et al. 2006; and unpublished Pacific Northwest data from Bary, Cogger, and Sullivan. See Additional Resources for citations.

^bCompost analyses are usually reported as P and K, while fertilizer labels use P₂O₅ and K₂O. To convert from P to P₂O₅, multiply by 2.3. To convert from K to K₂O, multiply by 1.2.

^cAvailable N is estimated percentage of total N that becomes available to plants during the first growing season after application.

^d“As-is” is typical dry matter content for compost at the time of sale or use.

^eBroiler litter sold as “compost” is often not fully composted because no water was added to facilitate microbial decomposition during the “composting” process.

Table 4. Other organic soil amendments: Typical nutrient content, dry matter, bulk density, and nitrogen availability

	N	P ^a	K ^a	Dry Matter	Density	Total N	C/N ratio	Available N ^b		
				lb/ton as-is ^c	lb/cubic yd ^d	% dry weight	% of total N	lb/ton as-is	lb/cubic yd	
Fresh shrimp waste	26	16	1	24	1400	5.4	5 ^e	75 to 100	20–26	14–19
Corvallis coffee grounds	12	1	3	30	1400	2.0	27	0 to 10	0–1	0–1
Uncomposted peppermint hay	18	3	14	30	1400	1.8	25	0 to 10 ^f	0–2	0–1
Composted peppermint hay	47	7	30	50	1400	4.7	8	5 to 10	2–5	1–4
Yard debris compost	18	3	8	55	900	1.7	17	0 to 20	0–4	0–1.6

^aOrganic amendment analyses are usually reported as P and K, while fertilizer labels use P₂O₅ and K₂O. To convert from P to P₂O₅, multiply by 2.3. To convert from K to K₂O, multiply by 1.2.

^bAvailable N is estimated percentage of total N that becomes available to plants during the first growing season after application.

^c“As-is” is typical dry matter content for yard debris compost. For other organic amendments, the dry matter values given are rough estimates based on limited data.

^dDensities listed as 1400 lb/cubic yd are estimates. Most products with high moisture (low dry matter) have a density close to 1400 lb/cubic yd. Use the dropped bucket method described in Worksheet B to get a more accurate estimate of bulk density.

^eShrimp waste contains carbon (C) in shells. The C:N value listed above is for shrimp meat (excludes C in shells).

^fPeppermint hay is the byproduct of distillation for mint oil, sometimes called a “slug” of peppermint. It immobilizes N for the first month after field application, but releases N thereafter. Cumulative N release for three years following application is about 20% of total N. For additional information, see the Nutrient Management Guide in Additional Resources.

Choosing a Laboratory

Choose a laboratory that routinely tests animal manure or other organic amendments and participates in a proficiency testing program. Proficiency testing programs are quality assurance and control programs where a third party submits samples and reviews results. See the Additional Resources section for information on choosing a laboratory and on proficiency testing programs.

Questions to ask when choosing a lab:

- Does the lab routinely analyze manure or other organic amendments?
- How many manure samples do they analyze in a year?
- Can they perform the tests you need?
- Are the results reported in the form you need?
- How are the samples handled before analysis? Do they thoroughly mix the samples? Do they dry the samples before analysis? (Drying samples prior to analyzing for ammonium nitrogen results in loss of most of the ammonium nitrogen from the sample.)
- Can the laboratory supply quality control information to you? Do they participate in a regional, national, or university proficiency testing program?

Laboratories report results on an as-is or a dry-weight basis. The as-is results are most helpful when determining application rates. The dry-weight results can be used to compare analyses over time and from different manure sources.

If you are not sure what reporting basis the lab is using, contact them for clarification. It is a good idea to ask for a sample reporting sheet before you choose a lab to make sure you can interpret their results (Figure 3).

To convert manure analyses reported on a dry-weight basis (in percentages) to an as-is basis (in lb/wet ton), multiply by 20 to convert the dry-weight percentage to lb/ton; then multiply by the decimal equivalent of the dry matter content.



Figure 3. Successful manure testing includes representative sampling, proper sample handling, and knowing how to interpret test results. If you have questions about laboratory reports, contact the lab for clarification. Photo by Andy Bary.

Example:

For beef manure at 23% solids and 2.4% nitrogen (N) on a dry-weight basis:

- Step 1. $4\% \times 20 = 48 \text{ lb N/ton dry weight}$
- Step 2. $48 \text{ lb N/ton dry weight} \times .23 = 11 \text{ lb N/ton as-is.}$

Sampling Manure

A nutrient analysis is only as good as the sample you take.

Samples must be fresh and representative of the manure.

Follow these steps carefully:

1. Ask the laboratory what type of containers they prefer. Also, make sure the laboratory knows when your sample is coming. Laboratories should receive samples within 48 hours of collection. Plan to collect and send your sample early in the week so the sample does not arrive at the lab on a Friday or a weekend.
2. If you have a bucket loader and a large amount of manure, use the loader to mix the manure before sampling.
3. Take 10 to 20 small samples from different parts and depths of the manure pile to form a composite sample. The composite sample should be about five gallons. The more heterogeneous your pile, the more samples you should take.
4. With a shovel or your gloved hands, thoroughly mix the composite sample. You may need to use your hands to ensure complete mixing. Wear waterproof gloves when mixing manure samples with your hands.
5. Collect about one quart of manure from the composite sample and place in an appropriate container.
6. Freeze the sample if you are mailing it. Use rapid delivery to ensure that it arrives at the laboratory within 24 to 48 hours. You can refrigerate the sample if you are delivering it directly to the lab.

See the Additional Resources section for more information on manure and compost sampling.

Published Values

Tables 2 and 3 show typical published values for livestock manure and compost. These values may not accurately represent your situation. The nutrient content of manure varies widely with the amount and type of bedding used, storage conditions, age of manure, manure handling, and animal diet. Nutrient values can vary by a factor of two or more from the values listed in the tables.

For example, leaving manure piles uncovered west of the Cascades will reduce manure nutrient content by leaching. The use of bedding reduces manure nutrients by dilution. The amount of reduction from either of these factors is difficult to determine without laboratory testing.

Analysis of horse manure samples collected from 37 farms in King County, Washington illustrates the variability in nitrogen concentration among manure sources (Figure 4). Only about one third of the samples with wood chip bedding had nitrogen concentrations similar to the mean of 1.3%, while the remainder had nitrogen concentrations less than 1% or greater than 1.5%. These differences would have a large effect on nitrogen availability, as described below. Dry matter also varied considerably among sources, ranging from 19% to 53%.




Figure 4. Range of nitrogen concentration in manure from 37 horse farms in King County, Washington.

Nitrogen Availability for Crop Growth

Manure application rates are usually nitrogen-based because nitrogen is usually the nutrient needed in the largest quantity for crop growth. Manure is not like commercial fertilizer in that it does not come with a guaranteed nitrogen availability. Nitrogen availability from manure varies greatly, depending on the type of animal, type and amount of bedding, and age and storage of manure.

Manure contains nitrogen in the organic and ammonium forms. The organic form releases nitrogen slowly, while ammonium-N is immediately available for crop growth. Ammonium-N can be lost to the atmosphere as ammonia gas when manure is applied to the soil surface. Most solid manures contain most of their nitrogen in the organic form, but poultry manure contains substantial ammonium-N.

Herbicides in Manure and Compost

Herbicides that are applied to forage and pasture crops can pass through animals into their manure, leaving a small amount of herbicide residue in the manure. In most cases, the levels of herbicide in manure are inconsequential and have no effect on the crops or soil where manure is applied.

A few herbicides that are active at very low concentrations and persist in the environment can cause problems in manure or manure compost. These include clopyralid and aminopyralid, herbicides that are used to kill broadleaf weeds in grass, grain, and some other crops. Residues from aminopyralid and clopyralid persist in manure and compost and have ruined sensitive crops, such as legumes (e.g., beans and peas), and nightshades (e.g., tomatoes, potatoes, peppers). The labels for these herbicides now stipulate that manure from animals eating treated forage should not be used on sensitive crops.

If you raise animals and plan on using, selling, or giving away manure or compost for vegetable production or home gardens, make sure that forages you grow or purchase have not been treated with these herbicides. If you obtain manure from a farmer or composter, ask them if they know that their forages are free of these materials. Refer to the Additional Resources section for more information on herbicides in manure and compost.

Poultry and other manures that contain a large proportion of ammonium-N may lose substantial nitrogen to the atmosphere if they are not tilled into the soil the same day they are spread. Ammonia loss is greater in warm, dry, and breezy conditions; less in cool, wet weather. Incorporating poultry manure immediately after incorporation with a harrow or disc is the best way to reduce ammonia loss. Irrigation immediately after application can also reduce ammonia loss.

There is no simple test to determine nitrogen availability for an individual manure sample. Use Tables 2 and 3 to estimate nitrogen availability from different manures and composts.

The nitrogen availability numbers in the Tables 2–5 are approximate ranges for each type of material. Use the lower part of the ranges if manure contains a large proportion of ammonium-N, such as poultry manure, and it is not tilled into the soil soon after application. Also, use the lower part of the ranges if the manure contains large amounts of bedding, or if the measured nitrogen content is lower than typical values. Use the higher part of the ranges if the manure contains less bedding or if the measured nitrogen content is higher than typical values.

Thorough composting generally reduces the rate of release of manure nitrogen by incorporating the nitrogen into more biologically resistant forms (Table 3). This may not be the case for material sold as composted poultry litter, however. Poultry litter (from broiler production) is a mixture of manure and wood shavings. Dry-stacked poultry litter will heat to high temperatures, but the composting process is often limited by insufficient moisture in the pile. Materials sold as “composted poultry litter” are frequently incompletely composted and have similar nitrogen availability as fresh poultry litter.

Incompletely composted poultry litter will have a strong ammonia odor and will have ammonium-N analyses above 2000 ppm (0.2% ammonium-N on a dry-weight basis).

Other Organic Fertilizers and Soil Builders

Other waste-derived organic amendments include recycled bulk agricultural and food processing wastes, such as peppermint hay, shrimp waste, and yard and food waste composts (Table 4). The bulk materials are similar to animal manure in that they can vary in nutrient concentration and handling characteristics. They may be either fresh or composted. Follow the guidelines for manure in this publication when choosing, analyzing, and applying these materials.

High nitrogen specialty products such as dried feather meal and fish meal are usually processed commercially to kill pathogens and render them into a form that is convenient to use (Table 5). They are typically sold as fertilizers, and their nutrient concentrations are on the product label. Use them as you would normally use fertilizers, calculating appropriate application rates and doing periodic soil testing. These materials can be applied using drop spreaders rather than manure spreaders.

Table 5. Organic-based specialty products^a: Typical nutrient content and nitrogen availability

	N	P ₂ O ₅	K ₂ O	Available N	
				% of total N	lb/ton
Feather meal	12	0	0	75–100	180–240
Blood meal	12	1	1	75–100	180–240
Fish meal	10	6	2	75–100	160–200
Fish emulsion	5	2	2	75–100	
Canola meal	5	1	1	40–60	40–60
Dried, pelletized poultry manure	4	3	3	40–60	30–50

^aAll of these products, except fish emulsion, have been dried to remove most moisture. Data sources include Gale et al. 2006; Sullivan et al. 2010; Darby et al. 2010, and product labels. These products are usually marketed commercially as fertilizers, and their labels show phosphorus and potassium concentrations in terms of P₂O₅ and K₂O. Note that this differs from the P and K analyses shown for manures and composts in Tables 2–4.

Horse manure or other manures with lots of woody bedding may temporarily reduce nitrogen availability rather than supply nitrogen for crop growth. Woody materials contain little nitrogen, and the microorganisms that break down the carbon in the wood will take available nitrogen from the soil to meet their nutrient needs. This process is called immobilization, and it reduces nitrogen availability to plants. Separated dairy manure solids are also low in nitrogen and can temporarily immobilize nitrogen after application to land. Expect at least a short-term nitrogen immobilization from manures and amendments containing less than 1.5% nitrogen.

If you have been applying manure to the same field for several years, residual organic nitrogen from past applications will contribute to the supply of nitrogen available to crops. You will need to reduce the manure application rate for fields that receive repeated manure applications.

Calculating the Application Rate

Now that you have determined the nutrient content and estimated the nitrogen availability of the manure you will be using, you are ready to determine the application rate. Application rates of manure are usually based on providing adequate nitrogen for crop growth. In most cases, supplying sufficient nitrogen will also provide adequate phosphorus and potassium for crop growth.

First, you will need to know the amount of nutrients your crop requires. Nutrient requirements for specific crops can be found in Extension production guides or from soil test recommendations.

Once you know the amount of nitrogen needed for your crop and the nutrient content of the manure, you can estimate the amount of manure to apply using a few simple calculations. See Worksheet C for help calculating application rates.

Observe the performance of your crops to help fine-tune your application rate. If you have a complicated nutrient management program, you may need to consult with a professional agronomist. See the Additional Resources section for a list of laboratories and professional agronomists, or contact your local Extension office.

Applying Manure

Manure Spreaders

The best of manure application estimates will not be useful if you don't know how much you're applying once you get to the field. You will need a spreader appropriate to the size of your farm, and you will need to calibrate it so you have confidence in your application rates.

Equipment

There are several ways to spread manure. For small amounts, you can use buckets or apply it from the back of a pickup truck. Using a mechanical manure spreader is easier and more effective for spreading large amounts of manure. Contracting with a commercial application service may be a good option for some farms.

There are three general types of solid manure spreaders: rear-delivery, side-delivery, and specialty spreaders. Rear-delivery spreaders work best with drier materials (Figure 5). Side-delivery spreaders handle both wet and dry materials. Specialty spreaders have the capability of spreading material onto narrow beds found in perennial crop production.



Figure 5. Applying manure compost before spring tillage using a rear-delivery spreader. Photo by Craig Cogger.

Calibrating Manure Applications

Calibrating your spreader will allow you to determine the actual application rate and adjust it to meet your target rate.

- You can weigh a spreader load and measure how much area the load covers (Figure 6). Use Worksheet A for calculations using this method.
- If you know the bulk density of the manure, you can calculate the volume of a spreader load rather than the weight. Use Worksheet B for calculations using this method.

Calibrating your spreader will allow you to convert loads of manure applied to tons per acre. The spreader load weight calibration is the most accurate method, but it requires truck scales to complete. Calibration improves accuracy of the overall application rate, but there will still be variability in the actual application across your field.

If you apply manure to a small area and use buckets or a pickup truck, it is still important to know how much you are applying. You can do this by weighing the buckets or determining the weight or volume of the pickup load.

Regardless of which method you use, recheck your measurements periodically to make sure you stay on target.

Timing Manure Applications

From a nutrient management perspective, the best time to apply manure to row crops is in the spring before planting. Fall application increases the risk of nutrient loss via leaching or runoff. Food safety and soil management impacts of manure application should also be considered.



Figure 6. Spreader load calibration method. Determine the area spread from the length and width of the band. Photo by Andy Bary.

Food safety requires a waiting period between manure application and crop harvest (see the section on Using Manure Safely). Using compost instead of manure provides greater flexibility in timing of application because compost has already been treated to reduce or eliminate human pathogens. Soil compaction risk during manure application is usually greater when soil is wet in the spring.

To reduce nutrient loss with fall manure application, apply manure early in the fall and plant a cover crop to help capture nutrients and prevent runoff. Alternatively, apply manure at moderate rates to pastures, hay fields, or other land with vegetated cover.

Using Manure Safely

Fresh manure sometimes contains pathogens that can cause diseases in humans. *Salmonella* and *E. coli* O157:H7 bacteria are among the most serious pathogens found in animal manure.

These pathogens are not taken up into plant tissue, but they can adhere to soil on plant roots or on the leaves or fruit of low-growing crops. Cooking destroys pathogens, but raw food carries a risk of pathogen exposure. The risk from pathogens is greatest for root crops (e.g., carrots and radishes) or leaf crops (e.g., lettuce or spinach), which have edible parts close to the soil. The risk is negligible for crops such as sweet corn, which does not have edible parts close to the soil, or for any crop that is cooked thoroughly.

Consider manure to be a potential source of pathogens. Pathogens die off naturally during extended storage or after field application. Die-off of bacterial pathogens occurs over days to months, depending on the pathogen and environmental conditions.

Composting manure at high temperatures will reduce pathogens, but you need careful quality control and daily temperature monitoring to make sure that all of the manure reaches conditions for pathogen kill (Figure 7). If you plan to compost manure, refer to the Additional Resources section for procedures to compost manure under conditions that destroy pathogens.

If you grow certified organic crops, you must follow the manure and compost practices in the National Organic Standards. These practices specify a waiting period of 120 days between the application of fresh manure and the harvest of high-risk crops and 90 days for low-risk crops. In practical terms, this means that many high-risk crops are not compatible with fresh manure being applied during the growing season in which the crops are grown.



Figure 7. If you compost manure on farm, monitor temperatures daily during active composting to document meeting time and temperature standards for pathogen kill. Photo by Andy Bary.

The National Organic Standards also specify procedures for composting manure, which include maintaining compost temperatures above 131°F for a minimum of three to 15 days, depending on the type of compost system used. Even if you don't grow certified organic crops, use the National Organic Standards as a minimum best practice when using animal manures. Refer to the National Organic Standards for more details.

The US Food and Drug Administration has drafted agricultural production regulations for protecting food safety, and they are developing research-based standards for manure and compost use. Refer to the Food Safety Modernization Act (FSMA) *Final Rule on Produce Safety* in Additional Resources for details on the food safety standards when manure is used.

Manure Storage

If you raise animals on your farm or import manure for crop production you may need to store manure on your farm prior to application. Proper manure storage conserves nutrients and protects surface water and groundwater. Storing manure can be as elaborate as keeping it under cover in a building or as simple as covering the manure pile with a tarp. The important point is keeping the pile covered and away from drainage areas and standing water. The storage location should also be convenient to your animals and crop production. To avoid inadvertent pathogen contamination of crops, choose a location with a low risk of runoff of manure onto cropland. If you decide to build a manure storage structure, refer to the Additional Resources section for a list of sources that cover this, or see your local Natural Resources Conservation Service office.

Using Soil Testing to Adjust Application Rates

Taking soil test samples and observing your crops can help you determine if manure application rates are adequate or if they need adjusting. The post-harvest soil nitrate test helps determine if you are applying too much manure. The post-harvest test measures nitrate-N remaining in the soil in the fall. If you apply too much manure, nitrate-N will accumulate in the soil, unused by the crop. When the fall and winter rains come, the nitrate will leach from the soil and become a potential contaminant in groundwater or surface water. Excess nitrogen can also harm some crops, delaying fruiting and increasing the risk of disease damage, winter injury, and wind damage. Although the post-harvest soil nitrate test outlined below was developed specifically for west of the Cascades, it also has use in nitrogen management in irrigated and dryland areas east of the Cascades.

To do a post-harvest soil nitrate test, sample the soil (zero to 12-inch depth) between August 15 and October 1. Timing is critical, particularly west of the Cascades. You want to sample after most crop uptake of nitrogen has occurred, but before the fall rains leach nitrate from the soil. Take a post-harvest nitrate sample as you would any other soil sample, collecting soil cores at multiple spots in the field and combining the cores into a composite sample. For details on soil sampling procedures, refer to Oregon State University Extension Publication EM 8832 *Post-harvest Soil Nitrate Testing for Manured Cropping Systems West of the Cascades* in Additional Resources. This publication also provides detailed background on the post-harvest test and interpretation.

If your post-harvest nitrate-N results are greater than 15–20 mg/kg, this suggests you are supplying more nitrogen than your crop needs, and you can reduce manure application rates. Post-harvest nitrate-N levels greater than 30 mg/kg are excessive.

When interpreting post-harvest soil nitrate results, consider the performance of your crop, too. If crop growth was poor because of drought, pests, or poor growing conditions, crop nitrogen uptake may have been less than expected, resulting in excess nitrate-N remaining in the soil profile even if manure applications were on target for a normal crop.

You can use routine soil tests to evaluate the soil for sufficiency or excess of other nutrients. A routine soil test includes phosphorus (P), potassium (K), calcium (Ca), magnesium (Mg), boron (B), and pH. West of the Cascades, a routine test should also include a lime requirement test. Soil tests east of Cascades should include electrical conductivity, a measure of soluble salts.

When soil test phosphorus and potassium are low, and you observe reduced crop growth, you can probably increase manure application rates. When soil test phosphorus and potassium are consistently excessive, you may need to decrease or eliminate manure applications, as described below. For additional information on soil tests for these and other nutrients, refer to Oregon State University Extension Publication EC 1478 *Soil Test Interpretation Guide* in Additional Resources.

Long-Term Effects of Manure Applications

Repeated applications of manure can increase soil organic matter content and the soil nutrient supply. Increasing organic matter often improves soil physical properties, such as bulk density, infiltration rate, and water-holding capacity. Because the supply of slow-release nutrients increases in soil, the amount of manure needed to meet crop needs will decline over time.

If you apply manure repeatedly to the same fields, it is important to have a regular soil testing program to track nutrient levels, as described in the Laboratory Analysis section of this publication. In drier regions (e.g., east of the Cascades), it is important to track soluble salts and cease applications of manure before salt accumulation reaches harmful levels. When manure applications are based on nitrogen need, manure usually supplies phosphorus and potassium in excess of crop needs. As a result, phosphorus and potassium can accumulate in soils with a history of manure applications, and may eventually reach excessive levels. Excess levels of soil phosphorus can increase the amount of phosphorus in runoff, increasing the risk of surface water degradation. Many crops can handle high levels of potassium, but blueberries and some other specialty crops can suffer from nutrient imbalances in soils with excessive potassium. Livestock can also be harmed by nutrient imbalances if they consume a diet of forages with high potassium levels.

If soil test phosphorus and potassium reach excessive levels, you may need to switch to a commercial fertilizer that supplies only nitrogen. If you are an organic farmer and do not use synthetic fertilizers, consider replacing some or all manure with legume cover crops or products such as feather meal with high nitrogen content and low phosphorus and potassium. Legumes will fix nitrogen from the atmosphere, but they do not supply phosphorus or potassium. Excess phosphorus and potassium levels can be reduced by growing forage crops, such as alfalfa, that will supply their own nitrogen but export considerable amounts of phosphorus and potassium in the dry matter that is removed. Reserve manure for fields that have lower phosphorus and potassium levels.

Worksheet A

Spreader Calibration Using a Full Spreader Load and Scales

Use this method to monitor the manure application rate on the entire field. After the initial weighing to determine the capacity of the application vehicle, the only tool you need is a field-measuring wheel or long tape.

1. Determine the weight of manure the spreader will hold using a truck scale to weigh the spreader when empty and full. Record the weights on Worksheet A.
2. Spread a load on the field using constant, even tractor speed and settings to cover the area uniformly. Spread in a rectangular pattern so the area calculation will be simple. Record tractor speed and gear settings used on Worksheet A.
3. Measure the length and width covered by one full load and compute the application rate in tons per acre using Worksheet A.
4. Adjust the application rate by changing tractor speed or gearing, or by making an adjustment on the manure spreader. After adjustment, you will need to repeat the calibration procedure until you have arrived at the desired application rate.
5. Keep the calibration records for future use.

Date _____

Field _____

Spreader ID _____

Operator _____

		Replicates				
		Example	1	2	3	4
Tractor gear		B2				
Tractor speed		2,000 rpm				
A	Weight of loaded spreader (lb)	12,000				
B	Weight of empty spreader (lb)	2,000				
C	Weight of manure in spreader (lb) [A – B]	10,000				
D	Length of area spread (ft)	1,300				
E	Width of area spread (ft)	25				
F	Area spread (sq ft) [D × E]	32,500				
G	Manure applied (lb/sq ft) [C ÷ F]	0.308				
H	Convert to ton/acre [G × 21.78]	6.7				

Note: Unshaded cells are information about manure and crop. Shaded cells are calculations you make.

Worksheet B

Spreader Calibration Using Manure Bulk Density

This method can be used if you know or estimate the manure bulk density, but it is not as accurate as other methods. You will need a tape measure, calculator, and a measurement or estimate of the manure bulk density.

1. Determine the length, width, depth, and stacking height of the manure spreader and enter the values in Worksheet B. Measure or estimate manure bulk density from Table 2 and enter it in the worksheet.
2. Calculate the volume and weight of manure in a spreader load using Worksheet B.
3. Spread a load on the field, using constant, even tractor speed and settings to cover field uniformly. Spread in a rectangular pattern so the area calculation will be simple. Record tractor speed and gear settings used on Worksheet B.
4. Measure the length and width covered by one full load and compute the application rate in tons per acre using Worksheet B.
5. Adjust the application rate by changing tractor speed or gearing, or making an adjustment on the manure spreader. After adjustment, you will need to repeat the calibration procedure until you have arrived at the desired application rate.

Date _____

Field _____

Spreader ID _____

Operator _____

		Replicates			
		Example	1	2	3
Tractor gear		B2			
Tractor speed		2,000 rpm			
A	Manure bulk density (lb/cubic yd.)	1,100			
B	Length of manure spreader box (ft)	7.0			
C	Width of manure spreader box (ft)	3.0			
D	Depth of manure to top of spreader box side boards (ft)	1.4			
E	Stacking height from the top of the spreader box side boards to top of pile (ft)	1.1			
F	Volume of manure in spreader (ft ³) [B × C × (D + (E × 0.8))]	48			
G	Weight of manure in the spreader (lb) [(A × F) ÷ 27]	1,951			
H	Length of application area covered by a full spreader load (ft)	245			
I	Width of area spread (ft)	4			
J	Area spread (sq ft) [H × I]	980			
K	Manure applied (lb/ sq ft) [G ÷ J]	1.99			
L	Convert to ton/acre [K × 21.78]	43			

Note: Unshaded cells are information about manure and crop. Shaded cells are calculations you make.

Bulk Density Estimate

You can estimate the bulk density of manure or compost using the “dropped bucket” method. You will need a 5-gallon bucket (make sure it is five gallons and not four) and a scale.

1. Set up the bucket by measuring 5 gallons of water into it and marking the 5-gallon level. Empty the bucket. Using a ruler, measure and mark 1/3 and 2/3 of the way to the 5-gallon level (Figure 8).
2. Weigh the empty bucket and record the weight.
3. Fill the bucket to the 1/3 mark with manure, and drop it 10 times from a height of one foot. The manure will settle to below the mark.
4. Add more manure to the 2/3 mark, and drop it 10 times again.
5. Add more manure to the full 5-gallon mark and drop it 10 times again.
6. Add enough manure to reach the 5-gallon mark again. **Do not drop the bucket this time.**
7. Weigh the full bucket and subtract the weight of the empty bucket. This is the weight of 5 gallons of manure.
8. Multiply this weight by 40 to calculate the bulk density in lb per cubic yard.
9. Repeat 2–3 times and take the average.



Figure 8. Calibrated bucket to use with dropped bucket method for estimating manure or compost bulk density. Photo by Andy Bary.

		<i>Example</i>	1	2	3
A	Weight of empty bucket (lb)	2.1			
B	Weight of full bucket (lb)	31.5			
C	Weight of manure (lb) [B – A]	29.4			
D	Bulk density (lb/cubic yd) [C × 40]	1176			

Note: Unshaded cells are information about manure and crop. Shaded cells are calculations you make.

Worksheet C

Calculating Manure Application Rates

Example: I am growing sweet corn and the nitrogen recommendation is 100 lb N per acre. I have broiler litter that has been tested by a laboratory and contains 50 lb N, 25 lb P, and 30 lb K per wet “as-is” ton of manure.

	Step	Units	Example	Your Value
A	Type of manure		<i>broiler litter</i>	
B	Crop		<i>sweet corn</i>	
C	Desired N application rate	lb N/acre	<i>100</i>	
D	Manure N concentration from laboratory analysis or Tables 2–5	lb N/ton as-is	<i>50</i>	
E	Phosphorus concentration from laboratory analysis or Tables 2–5	lb P/ton as-is	<i>25</i>	
F	Potassium concentration from laboratory analysis or Tables 2–5	lb K/ton as-is	<i>30</i>	
G	Plant availability of N from Tables 2–5	%	<i>50</i>	
H	Calculate manure available N [$D \times (G \div 100)$]	lb N/ton as-is	<i>25</i>	
I	Calculate application rate [$C \div H$]	ton manure/acre as-is	<i>4</i>	
J	Calculate amount of phosphorus applied [$I \times E \times 2.3$]	lb P ₂ O ₅ /acre	<i>230</i>	
K	Calculate amount of potassium applied [$I \times F \times 1.2$]	lb K ₂ O/acre	<i>144</i>	

Note: Unshaded cells are information about manure and crop. Shaded cells are calculations you make.

Additional Resources

Source Data for Manure and Other Amendment Nutrient Concentrations and Availability

Bary, A. 2010. [Horse Manure and Soil Nitrogen](#).

Brown, C. 2013. [Available Nutrients and Value for Manure from Various Livestock Types](#). Ontario Ministry of Agriculture and Food publication 13-043.

Darby, H.K. Hills, E. Cummings, and R. Madden. 2010. [Assessing the Value of Oilseed Meals for Soil Fertility and Weed Suppression](#). University of Vermont Extension.

Gale, E.S., D.M. Sullivan, C.G. Cogger, A.I. Bary, D.D. Hemphill, and E.A. Myhre. 2006. Estimating Plant Available Nitrogen Release from Manures, Composts, and Specialty Products. *Journal of Environmental Quality* 35: 2321–2332.

Laboski, C.A.M. and J. B. Peters. 2012. [Nutrient Application Guidelines for Field, Vegetable, and Fruit Crops in Wisconsin](#). *University of Wisconsin Extension Publication A2809*. University of Wisconsin.

Larney, F.J., D.M. Sullivan, K. E. Buckley, and B. Eghball. 2006. The Role of Composting in Recycling Manure Nutrients. *Canadian Journal of Soil Science* 86: 597–611.

Sullivan, D.M., N.A. Andrews, J.M. Luna, and J.P. McQueen. 2010. [Estimating N Contribution from Organic Fertilizers and Cover Crop Residues Using Online Calculators](#). World Congress of Soil Science.

Kusonwiriya Wong, C., D.M. Sullivan, D.D. Hemphill, C.G. Cogger, A.I. Bary, and L. Myhre. 2014. Predicting Plant-Available Nitrogen from Organic Amendment in the Second Year After Application. *Acta Horticulture* 1018: 325–333.

Manure Handling and Composting on the Farm

Bary, A., G. Hoogenboom, and S. Hill. 2015. [Compost Calculator Phone App](#). Washington State University.

Brewer, L., N. Andrews, D. Sullivan, and W. Gehr. 2013. [Agricultural Composting and Water Quality](#). *Oregon State University Extension Publication EM 9053*. Oregon State University.

Martin, G. and D. Beegle. 2008. [Manure Spreader Calibration](#). Agronomy Facts 68. Penn State University Extension.

Midwest Plan Service. 2001. [Manure Storages](#). Iowa State University MWPS-18-S2.

Rynk, R. 1992. [On-Farm Composting Handbook](#). Northeast Regional Agricultural Engineering Service Publication NRAES-54.

Sullivan, D.M. 2008. [Estimating Plant-available Nitrogen from Manure](#). *Oregon State University Extension Publication EM 8954-E*. Oregon State University.

Herbicides in Manure and Compost

[Clopyralid in compost](#). Washington State University Puyallup Research and Extension Center.

Davis, J., S.E. Johnson, and K. Jennings. 2015. [Herbicide Carryover in Hay, Manure, Compost, and Grass Clippings](#). North Carolina Cooperative Extension. AG-727W.

Food Safety

US Food and Drug Administration. 2015. [FSMA Final Rule on Produce Safety](#).

Lists of Laboratories and Consultants

The first three resources list only labs that participate in proficiency testing quality control programs.

Minnesota Department of Agriculture. [Certified Manure Testing Laboratories](#).

[North American Proficiency Testing Program](#).

US Composting Council. [Compost Analysis Proficiency Testing Program](#).

Washington State Pest Management Resource Service. [Analytical Laboratories and Consultants Serving Agriculture in the Pacific Northwest](#). Washington State University.

Organic Nutrient Management and Soil Testing

Collins, D.C. Miles, C. Cogger, and R. Koenig. 2013. [Soil Fertility in Organic Systems: A Guide for Gardeners and Small Acreage Farmers](#). *Washington State University Extension Publication PNW646*. Washington State University.

Fery, M. and E. Murphy. [2013 A Guide to Collecting Soil Samples for Farms and Gardens](#). *Oregon State University Extension Publication EC 628*. Oregon State University.

Hart, J.M., D.M. Sullivan, M.E. Mellbye, A.G. Hulting, N.W. Christensen, and G.A. Gingrich. 2010. [Nutrient Management Guide: Peppermint \(Western Oregon\)](#). *Oregon State University Extension Publication EM 9018-E*. Oregon State University.

Horneck, D.A., D.M. Sullivan, J. Owen, and J.M. Hart. 2011. [Soil Test Interpretation Guide](#). *Oregon State University Extension Publication EC 1478*. Oregon State University.

Moore, A., M. de Haro-Marti, and L. Chen. 2015. [Sampling Dairy Manure and Compost for Nutrient Analysis](#). *University of Idaho Extension Publication PNW 673*. University of Idaho.

Sullivan, D.M. and C.G. Cogger. 2003. [Post-harvest Soil Nitrate Testing for Manured Cropping Systems West of the Cascades](#). *Oregon State University Extension Publication EM 8832*. Oregon State University.

Using Organic Amendments

Bell, N., D. Sullivan, and T. Cook. 2009. [Mulching Woody Ornamentals with Organic Materials](#). *Oregon State University Extension Publication EM 1629*. Oregon State University.

Bell, N., D.M. Sullivan, L.J. Brewer, and J. Hart. 2003. [Improving Garden Soils with Organic Matter](#). *Oregon State University Extension Publication EC 1561*. Oregon State University.

Sullivan, D.M., R. Costello, D.R. Bryla, and B.C. Strik. 2015. Organic Soil Amendments and Mulches for Blueberry: The Good, the Bad, and the Ugly. *Oregon State University Blueberry School 2015*: 46–52.

National Organic Standards

USDA. [National Organic Standards](#).

Published and distributed in furtherance of the Acts of Congress of May 8 and June 30, 1914, by Washington State University Extension, Oregon State University Extension Service, University of Idaho Extension, and the U.S. Department of Agriculture cooperating. WSU Extension programs, activities, materials, and policies comply with federal and state laws and regulations on nondiscrimination regarding race, sex, religion, age, color, creed, and national or ethnic origin; physical, mental, or sensory disability; marital status or sexual orientation; and status as a Vietnam-era or disabled veteran. Washington State University Extension, Oregon State University Extension Service, and University of Idaho Extension are Equal Opportunity Employers. Evidence of noncompliance may be reported through your local Extension office. Trade names have been used to simplify information; no endorsement is intended.

Pacific Northwest Extension publications contain material written and produced for public distribution. You may reprint written material, provided you do not use it to endorse a commercial product. Please reference by title and credit Pacific Northwest Extension publications.

Copyright 2016 Washington State University

Order Information:

Washington State University Extension http://pubs.wsu.edu Fax 509-335-3006 Toll-free phone 800-723-1763 ext.pubs@wsu.edu	Oregon State University Extension Service http://extension.oregonstate.edu/catalog Fax 541-737-0817 Toll-free phone 800-561-6719 puborders@oregonstate.edu	University of Idaho Extension http://www.cals.uidaho.edu/edComm/catalog Fax 208-885-4648 Phone 208-885-7982 calspubs@uidaho.edu
--	--	---

Pacific Northwest Extension publications are produced cooperatively by the three Pacific Northwest land-grant universities: Washington State University, Oregon State University, and the University of Idaho. Similar crops, climate, and topography create a natural geographic unit that crosses state lines. Since 1949, the PNW program has published more than 650 titles, preventing duplication of effort, broadening the availability of faculty specialists, and substantially reducing costs for the participating states. Published May 2016.